

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 864 560 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:

16.09.1998 Bulletin 1998/38

(51) Int. Cl.⁶: **C07C 235/30, C07D 309/30**

(21) Application number: 97107059.4

(22) Date of filing: 29.04.1997

(84) Designated Contracting States:

AT BE CH DE DK ES FI FR GB GR IE IT LI LU MC
NL PT SE

(30) Priority: 13.03.1997 US 816574

(71) Applicant:

RANBAXY LABORATORIES, LTD.
New Delhi 110 019 (IN)

(72) Inventors:

- Khanna, Jag Mohan
New Delhi 110 049 (IN)

• Kumar, Yatendra

Gurgaon 122001, Haryana (IN)

• Thaper, Rajesh Kumar

Haryana (IN)

• Misra, Satyananda

Janakpuri, New Delhi 110 058 (IN)

• Kumar, S.M. Dileep

Kalkaji Extension, New Delhi 100 019 (IN)

(74) Representative:

Cohausz & Florack

Patentanwälte

Kanzlerstrasse 8a

40472 Düsseldorf (DE)

(54) **Key intermediates in the manufacture of simvastatin**

(57) A process for preparing simvastatin from lovastatin or mevinolinic acid in salt form comprises treating either starting material with cyclopropyl or butyl amide, the pyranone ring thereby being opened when lovastatin is the starting material, adding a methyl group to the 2-methylbutyrate side chain, and thereafter closing the open pyranone ring to produce simvastatin. The process is performed without protecting and deprotecting the two hydroxy groups of the open pyranone ring. In a preferred embodiment, the starting material is treated with cyclopropyl amine which produces simvastatin via the novel intermediate lovastatin cyclopropyl amide.


EP 0 864 560 A1

Description**Reference to Related Application**

This patent application is related to a patent application entitled "PROCESS FOR MANUFACTURING SIMVASTATIN FROM LOVASTATIN OR MEVINOLINIC ACID" filed concurrently herewith.

Background of the Invention

Compounds of structure I shown below are very active antihypercholesterolemic agents that function to limit cholesterol biosynthesis by inhibiting the enzyme HMG-CoA reductase.



Compounds of structure I include the natural fermentation products mevinolin (structure Ia where R=CH₃, disclosed in U.S. Pat. No. 4,231,938, and also known as lovastatin), compactin (structure Ib where R=H, disclosed in U.S. Pat. No. 3,983,140), and a variety of semi-synthetic and totally synthetic analogs thereof, all having the natural 2-methylbutyrate side chain.

Compounds of structure II shown below having a 2,2-dimethylbutyrate side chain (e.g., simvastatin, structure IIa where R=CH₃) are known to be more active inhibitors of HMG-CoA reductase than their 2-methylbutyrate analogs, and thus of greater utility in the treatment of atherosclerosis, hyperlipemia, familial hypercholesterolemia, and similar disorders.



II

a: R = CH₃

15

20

25

30

- (1) de-esterification of the 2-methylbutyrate side chain;
- (2) protection of the 4-hydroxy group of the pyranone ring;
- (3) re-esterification of the side chain to form the desired 2,2-dimethylbutyrate; and
- (4) deprotection of the 4-hydroxy group.

35


40

45

Summary of the Invention

55

REACTION SCHEME



Salient features of the invention include:

(a) Simvastatin may be prepared from a salt of mevinolinic acid as the starting material rather than lovastatin.

(b) Simvastatin is prepared in a four step process which does not include the protection and deprotection of the two hydroxy groups of the open pyranone ring. Heretofore, the methods described in the prior art require the protection and deprotection of the hydroxy groups as essential steps for the preparation of simvastatin.

(c) In a preferred embodiment, the novel intermediate lovastatin cyclopropyl amide (IIIb) is prepared from the starting material (lovastatin or a salt form of mevinolinic acid). This novel intermediate is then converted into a second novel intermediate (IVb).

Simvastatin prepared by the inventive process has certain advantages in commercial manufacture. The purity and yield of simvastatin from the inventive process is high, with less consumption of reagents, time, labor, and cost, and in fewer steps.

Detailed Description of the Invention

The starting material for the method of the present invention is lovastatin (Ia) or mevinolinic acid (Ic), the latter being the open ring form of lovastatin. Mevinolinic acid is produced by fermentation of *Aspergillus terreus* (see U.S. Patent No. 4,231,938). Either lovastatin (Ia) or the salt form of mevinolinic acid (Ic) will typically be employed as the starting material. The term "mevinolinic acid" unless otherwise specified includes suitable salt forms thereof; any salt which will not interfere with other reagents or conditions used in carrying out the present invention is permissible. The alkali metal salts, e.g., Na and K, or, preferably, the ammonium salt form, may be employed.

Lovastatin is obtained from mevinolinic acid via lactonization by methods known in the art, and is isolated by loss incurring crystallization techniques. There is ca. 20% material loss in converting naturally occurring mevinolinic acid to lovastatin. One objective of this invention is to eliminate this loss by employing mevinolinic acid as the starting material rather than lovastatin.

According to the novel process of the present invention, simvastatin (IIa) is prepared by reacting lovastatin (Ia) or mevinolinic acid (Ic), preferably in the ammonium salt form, with an n-alkylamine or a cycloalkylamine of formula R_3-NH_2 , wherein R_3 is C_3-C_6 . Preferably, the amine R_3-NH_2 is cyclopropylamine and the intermediate that is formed is lovastatin cyclopropylamide (IIIb). Alternatively, the amine R_3-NH_2 is n-butylamine and the intermediate that is formed is lovastatin n-butylamide (IIIa).

The cyclic intermediate amide IIIa or IIIb thus prepared is dissolved in dry tetrahydrofuran and is added to a solution of alkali metal amide, e.g., lithium pyrrolidide in THF at a temperature of about $-35^{\circ}C$ to $-40^{\circ}C$. Lithium pyrrolidide is generated *in situ* by the reaction of n-BuLi with pyrrolidine in THF. The solution of intermediate amide IIIa or IIIb, and the *in situ* generated base is aged at about -35° to $-40^{\circ}C$ for 1 hour, and dried alkyl halide, preferably methyl iodide (2 to 3.5 mol. eq.), is added in a single charge. The contents are stirred at about $-30^{\circ}C$ for 1 hour, warmed to $-10^{\circ}C$ and aged at this temperature for 20 minutes. Water is added to the reaction mixture and the layers are separated. The THF layer is washed with a mineral acid, preferably hydrochloric acid, and then concentrated to an oily mass which contains the intermediate IVa or IVb.

To the solution of intermediate IVa or IVb in methanol, without purification, is added 2.0N NaOH and is refluxed at about 80° to $81^{\circ}C$ for 2 to 6 hours, preferably for 2 hours. The mixture is cooled to about $50^{\circ}C$, and the methanol is stripped off at reduced pressure and then diluted with water. The mixture is then acidified at about $10^{\circ}C$ by careful addition of 2.0N HCl maintaining a pH of 6, extracted with ethyl acetate, while further acidifying to a pH of 4. The ethyl acetate layer is separated and the hydrolyzed product is precipitated out as a crystalline material in the form of ammonium salt (V) by slow addition of methanolic ammonium hydroxide over a period of 30 minutes at about 22° - $25^{\circ}C$, followed by cooling to $5^{\circ}C$.

The ammonium salt (V) is relactonized by heating in a hydrocarbon solvent such as toluene. The mixture is suspended in toluene, heated and stirred at about 100° - $110^{\circ}C$ for 2 to 10 hours, preferably at about 105° for 5 hours, under a sweep of nitrogen gas. The mixture is then cooled to about $35^{\circ}C$, carbon treated, filtered, and the filtrate is concentrated under reduced pressure at about a $60^{\circ}C$ bath temperature to one-tenth of its original volume. The lactone is crystallized from a hydrocarbon solvent such as cyclohexane to give simvastatin (IIa) of high purity.

In a most preferred embodiment of the novel process of this invention, mevinolinic acid (Ic) in its ammonium salt form is used as the starting material and is converted into lovastatin cyclopropylamide (IIIb). The mevinolinic acid salt is suspended in toluene and refluxed for 5-6 hours at about 100° - $110^{\circ}C$, preferably at about 105° - $107^{\circ}C$. The resulting solution is concentrated to one tenth of its volume by distilling the toluene at reduced pressure at about 50° - $55^{\circ}C$ bath temperature. Cyclopropylamine is added at $30^{\circ}C$ and the mixture again heated for 4-5 hours at about 40° - $50^{\circ}C$. Toluene and unreacted cyclopropylamine are distilled off at reduced pressure to afford lovastatin cyclopropylamide (IIIb) in quantitative yield. The cyclic amide thus prepared is used as such without purification in the next step of C-methylation in a similar way as described above without the protection of the dihydroxy system to give simvastatin (IIa).

The following examples further illustrate the present invention:

Example I**Preparation of Simvastatin (IIa) from Mevinolinic Acid Ammonium Salt (Ib) Using Cyclopropylamine**

Step-1 : N-Cyclopropyl-7-[1,2,6,7,8,8a(R)-hexahydro-2(S), 6(R)-dimethyl-8(S)-[[2(S)-methylbutanoyl]oxy]-1(S)-naphthyl]-3(R),5(R)-dihydroxyheptanoic acid amide (IIIb)

Mevinolinic acid ammonium salt (Ic) (12.5 g, 0.296 moles) was suspended in toluene (400 ml). The mixture was heated and stirred at 105-107° under a sweep of nitrogen gas for 5 hours. The temperature was lowered to 60°C and about 350ml of toluene was distilled off. Cyclopropylamine (12 ml, 0.172 moles) was added at 30°C and the solution was again stirred at 40-45°C for 4 hours. The toluene was slowly stripped off at reduced pressure and at a bath temperature of 55°C to afford the title compound in gum-like form. HPLC purity = 99.63%; ¹H NMR (CDCl₃, 300 MHz): δ 0.495 (m, 2H), δ 0.50 (m, 2H), δ 0.86 (m, 6H), δ 1.08 (m, 6H), δ 2.3 (d, 2H), δ 2.6 (m, 1H), δ 3.7 (m, 1H), δ 4.18 (m, 1H), δ 5.4 (m, 1H), δ 5.5 (bt, J = 3.0 Hz, 1H), δ 5.7 (dd, J = 6.1, 9.5 Hz, 1H), δ 5.9 (d, J = 9.6 Hz, 1H), δ 6.2 (bt, J = 5.3 Hz, 1H); IR(CHCl₃): λ max 3500 - 3300 (b), 3000, 1740, 1660, 1530, 1450, 1210, 860, 760 cm⁻¹.

This gum was directly used in the next step without purification.

Step-2 : N-Cyclopropyl-7-[1,2,6,7,8,8a(R)-hexahydro-2(S), 6(R)-dimethyl-8(S)-[[2,2-dimethyl-butanoyl]oxy]-1(S)-naphthyl]-3(R),5(R)-dihydroxyheptanoic acid amide (IVb)

Pyrrolidine (13.5 ml, 0.163 moles) in THF (50 ml) was cooled to -45°C and n-butyl lithium in hexane (1.6 M, 100 ml, 0.163 moles) was added under nitrogen at such a rate so as to maintain a temperature of -20° to -15°C. The mixture was stirred at -20° to -25°C for 40 minutes after the completion of addition.

A solution of compound IIIb in THF (300 ml) was prepared as described above and then slowly added via a cannula maintaining the temperature below -35°C during the whole addition. The solution was aged at -35° to -40°C for 1 hour. Sieve dried methyl iodide (4.82 ml, 0.077 moles) was added in a single lot. The off-white, cloudy solution thus obtained was stirred for 1 hour at -35° to -33°C, then warmed to -10°C and aged for 20 minutes. Distilled water (105 ml) was added to the reaction mixture and the contents vigorously stirred for 5 minutes. The layers were separated and the upper THF layer was treated with 1N HCl (105 ml). The THF layer was concentrated at reduced pressure to a volume of about 40 ml to yield the title compound IVb, wherein R₃ is cyclopropyl.

Step-3 : 6(R)-[2-[8(S)-(2,2-dimethylbutyryloxy)-2(S),6(R)-dimethyl-1,2,6,7,8,8a(R)-hexahydro-1(S)-naphthyl]ethyl]-4(R)-hydroxy-3,4,5,6-tetrahydro-2H-pyran-2-one (IIa) (Simvastatin)

To the concentrated solution from the previous step containing compound IVb was added an aqueous solution of NaOH (2N, 25 ml) and MeOH (175 ml). The mixture was refluxed at 80°-81°C for 2 hours. The mixture was cooled to 50°C, maximum MeOH was stripped off at reduced pressure, and diluted with water (90 ml). Acidification of the solution was done at 10°C by careful addition of 2N HCl (pH 6). Ethyl acetate (200 ml) was added and the mixture was vigorously agitated while further acidifying to pH of 4. The ethyl acetate layer was separated from the aqueous one. A solution of NH₄OH and MeOH (1:1, 10 ml) was slowly added in over 30 minutes at 22°-25°C. The precipitates were stirred at 25°C for 1.5 hour, cooled to 5°C and stirred at this temperature for 30 minutes. Filtration followed by washing with cold ethyl acetate (25 ml) and drying in vacuum at 35°C afforded the ammonium salt (V).

The crude ammonium salt (V) from the previous step (10 g, 0.022 moles) was suspended in toluene (350 ml). The mixture was heated and stirred at 105°C under a sweep of nitrogen for 5 hours. The solution was cooled at 35°C, activated charcoal (0.5 g) was added, stirred for 0.35 hour, and then filtered through celite bed. The filtrate was concentrated in vacuum to a volume of 40 ml at 60°C bath temperature. Cyclohexane (125 ml) was added and the solution again refluxed for 15 minutes, cooled for 1 hour to 25°C, and further cooled to 10° - 12°C for 30 minutes. The precipitates were stirred at 10° - 12°C for 30 minutes, filtered and washed with cold cyclohexane (50 ml), dried in vacuum at 35°C to give white crystalline product (IIa) which was further crystallized from absolute ethanol to afford the title product of >99% purity.

Example II**Preparation of Simvastatin (IIa) from Lovastatin (Ia) Using Cyclopropylamine**

Step-1 : N-Cyclopropyl-7-[1,2,6,7,8,8a(R)-hexahydro-2(S), 6(R)-dimethyl-8(S)-[2(S)-methylbutanoyl]oxy]-1(S)-naphthyl]-3(R),5(R)-dihydroxyheptanoic acid amide (IIIb)

Lovastatin (Ia) (12.5 g, 0.03 moles) was suspended in cyclopropylamine (13 ml, 0.174 moles) at 25°C. The mixture was slowly heated to 40-45°C and stirred at this temperature for 5 hours. Excess amine was evaporated off at reduced pressure and a 40°C bath temperature to afford the title compound in gum-like form. This gum was directly used in the next step without purification.

Steps 2-3: Compound (IIIb) obtained from the previous step was converted into simvastatin (IIa) by following the same procedure as described for Example I.

Examples III and IV


Preparation of Simvastatin (IIa) from Lovastatin (Ia) and Mevinolinic Acid Ammonium Salt (Ic) Using n-Butylamine

Simvastatin (IIa) was prepared from lovastatin (Ia) and mevinolinic acid ammonium salt (Ic) as starting materials following the same steps as described in Examples I and II but substituting equimolar quantities of n-butylamine in place of cyclopropylamine.

While the invention has been described by reference to specific embodiments, this was for purposes of illustration only. Numerous alternative embodiments will be apparent to those skilled in the art and are considered to be within the scope of the claims.


Claims

1. A compound of structural formula III':




wherein R' is methyl, R₃ is cyclopropyl, and wherein n=1 or 2.


2. The compound of claim 1 wherein $n = 1$.
3. The compound of claim 1 wherein $n = 2$.
4. A process for preparing a compound of structural formula III:



wherein R_3 is cyclopropyl, comprising treating a compound of structural formula I':




wherein R is:



and R_2 is Na, K, or NH_4 , with cyclopropyl amide.

5. The process of claim 4 wherein R_2 is NH_4 .

6. The process of claim 4 further comprising methylating the methylbutyrate side chain to produce a compound of structural formula IV:



wherein R_3 is as defined above.



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 97 10 7059

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
A,D	US 3 983 140 A (ENDO AKIRA ET AL) 28 September 1976 * claims *	1-6	C07C235/30 C07D309/30
A,D	US 4 231 938 A (MONAGHAN RICHARD L ET AL) 4 November 1980 * claims *	1-6	
A	CLAREMON D A ET AL: "AN EFFICIENT CHEMOSELECTIVE SYNTHESIS OF NITRILES FROM PRIMARY AMIDES" TETRAHEDRON LETTERS, vol. 29, no. 18, 1988, pages 2155-2158, XP000676446 * the whole document *	1-6	
A	ASKIN D ET AL: "SYNTHESIS OF SYNVINOLIN: EXTREMELY HIGH CONVERSION ALKYLATION OF ANESTER ENOLATE" JOURNAL OF ORGANIC CHEMISTRY, vol. 56, no. 16, 2 August 1991, pages 4929-4932, XP000676448 * the whole document *	1-6	
A	US 4 908 452 A (CLAREMON DAVID A) 13 March 1990 * claims *	1-6	
A	US 5 393 893 A (KUBELA RUDOLF ET AL) 28 February 1995 * claims *	1-6	
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 20 April 1998	Examiner Sánchez García, J.M.
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

EPO FORM 1503 03.82 (P04C01)